

Global mortality rate and statistical results of Coronavirus

S. Saeed, A.S. Hashim, H. Khan

Neurospinal and Cancer Care Institute and Hospital, Karachi, Pakistan

ABSTRACT:

- **Objective:** The existing facts on the traits of the unconventional coronavirus disease (COVID-19) Computerized Tomography (CT) scan with statistical results of tracking database are sent in specific journals, and a constant evaluation of the literature has not yet been compiled.
- **Patients and methods:** COVID-19 disease, a systemic inflammation that impacts several organs, may be defined as a sickness that affects all age degrees, has a high mortality price, extreme negative results, and a chief economic burden. The Embase internet site, PubMed, OVID Medline, Cochrane Library, Science Direct, and Wiley became used to conduct a scientific assessment of these studies. COVID-19 patients' physiological acoustics are crucial for isolating them from other respiration infections. This report seeks to provide a summary of the Coronavirus epidemic, which has had a sizeable impact on humans' lives everywhere in the world. Plenty of humans have died, and the worldwide death rate from network-obtained pneumonia has reached three hundred million and continues to acquire pneumonia.
- **Results:** The most critical reason for shifting COVID-19 sufferers to the in-depth care unit is acute respiratory misery syndrome, which is likewise the leading motive of loss of life in these patients. COVID-19 sickness is thought to affect the blood vessels, lungs, heart, anxious gadget, liver, digestive system, kidneys, pores and skin, and other organs. Adults, infants, newborns, pregnant women, and the elderly are all infected, and it is spread through the air and droplets.
- **Conclusions:** This systematic analysis of the latest literature on COVID-19 includes detail on the CT disease's baseline and follow-up characteristics.
- **Keywords:** COVID-19, Patients, CT scan, Disease, Literature.

INTRODUCTION

The new Coronavirus (SARS-CoV-2) epidemic commenced in overdue December 2019 and soon spreads across the world, wreaking havoc on public health systems. Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) causes respiratory disorders (COVID-19), which can range from asymptomatic pinnacle respiratory contamination to acute respiration distress syndrome (ARDS), a couple of organ failure, and death¹. This accelerating disaster has inspired professionals all around the international

to step up their tries to come to be privy to inexperienced and comfy COVID-19 care and preclusion solutions. On January 30, 2020, the World Health Organization (WHO) Emergency Committee declared the COVID-19 epidemic a global fitness emergency of global importance (ESPII) and issued a joint studies timetable of eight precise regions to be prioritized in the course of the pandemic's first duration^{2,3}. The first wave of published facts for therapeutic modalities targeted largely on retrospective research and indirect findings from comparable fields⁴. Countries and regulators have already modified rapid pathways



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-nc-sa/4.0/)

for moral and other permissions^{5,6}, and courses have now increased reporting, each of which allows for the quick conduct and dissemination of randomized clinical trials (RCTs) to ensure a ‘2d wave of best records with enough assessment of advantages and harms (preprint)^{7,8}. The World Health Organization has created a global technology roadmap domestic page to help and music the selection of key research dreams and their proper execution⁹. While many versions of trial registry databases are to be had for download online¹⁰⁻¹², researchers can war to gain a summary of the various contemporary, expected, and constantly converting trials. As a result, we are hoping to offer a direct précis of the cures and core methodological functions of the proposed randomized medical trials at the preclusion and cure of COVID-19 with statistical results through regression test conducted by SPSS software through tracking in overall the world. Recent findings of a complementary thrombotic microangiopathy-like syndrome in COVID-19 have pressured the disease’s pathophysiology. The intention of this examination became to do a scientific review of previously published systematic evaluations to offer complete and up-to-date statistics on COVID-19.

MORTALITY RATE OF COVID-19 WORLDWIDE

The columns in the desk beneath display the number of deaths according to one hundred said cases (the case-monitored mortality rate) or according to a 100,000 population for the 20 international locations currently worst stricken by COVID-19 worldwide (this represents the general population of a rustic, with both Confirmed instances and wholesome human beings). As seen in Figure 1, the countries with the largest number of deaths have the highest number of deaths in proportion to COVID-19 cases or population, not exactly the highest number of deaths overall. This research is conducted by the tracking system of software to analyze the COVID-19 cases including confirmed, reported and death cases of COVID-19.

Figure 1 shows the fatality ratio of death cases of COVID-19 patients all over the world. The diagonal lines in the above Figure 1 reflect various case fatality rates (no. of deaths divided by the number of confirmed cases). The highest mortality rates have been recorded in countries at the top of the rankings. The 20 countries with the largest no. of expiries (death) from COVID-19 are denoted by points with black borders. However, we can see the name of the country and the percentage value. Toggle between the following options by using the boxes at the top: 1) Deaths as a percentage of the total number of reported cases in the country; Figure 2 portrays the number of deaths per 100,000 inhabitants (this reflects the general population of a region, including reported cases and safe people).

Figure 2 shows the confirmed death cases of COVID-19 patients with graphical representation. Here are some systematic reviews mentioned in this research which declares the recent condition of COVID-19 patient cases around the world which is given below.

A Systematic Review of Coronavirus Pandemic: R1: What is the overall communal death rate for pandemic coronavirus patients?

Coronavirus is fundamental contamination including various organs, influencing all age gatherings, with a high death rate, genuine unfavorable results, and a critical monetary weight that should be depicted. The author used PubMed, OVID Medline, the Cochrane Library, N Engl, J Med, AHA, and LITCOVID publishing bodies to investigate the current information of COVID-19. The sureness of the proof was evaluated utilizing the Grade approach. Meta-investigation was performed by an irregular impacts model. 71 qualified deliberate surveys were remembered for the examination. 86.5% of them are of high caliber and 13.5% of them are mid-range quality. The aftereffects of the meta-examination are introduced in even structure and the excess outcomes are introduced in account structure coronavirus incorporates the veins, lungs, heart, sensory sys-




Figure 1. Fatality ratios of overall death cases.




Figure 2. Confirmed death cases.

tem, liver, stomach related framework, kidneys, eyes, and different organs and influences grown-ups, kids, infants, pregnant ladies and the older, and is communicated through the air and by beads. The comorbidities related to coronavirus are HBP 20.7%, CVD 9.6%, DM 9.55%, respiratory sickness 7% and 9% smoking. The prognostic variables for passing among coronavirus cases are intense heart injury, cardiovascular infection, diabetes, respiratory illness and hypertension. The variables prescient of sickness seriousness are CVD and HT. Fever, windedness, and smoking were prescient variables for illness movement. There is no particular antiviral cure. Preventive processes have been suggested including an actual distance of 2 meters and that is only the tip of the iceberg, utilization of individual defensive hardware, keeping away from parties, isolate and disconnection. Telemedicine, online training, and homeschooling are strongly encouraged. The vaccine is on its way to emerging and there are concerns about a highly effective vaccine. Modulation of cardiovascular risk and cardiac metabolism has become a cornerstone of the sustainable control of the pandemic¹.

R2: What Are the Effects of Deficiency of Vitamin D for Pandemic Coronavirus?

There are a few investigations connecting nutrient D inadequacy to the danger of more awful results in patients with coronavirus (COVID-19). Our point was to direct an efficient survey of the current writing on the job of nutrient D lack in coronavirus contamination and mortality and whether supplementation with high dosages of nutrient D could be useful in diminishing dangers and improving results. A precise hunt of PubMed, EMBASE and the Cochrane Library was performed until June 5, 2020. The nature of the included examinations was surveyed utilizing the Downs and Black predisposition hazard scale. The accessible writing was assessed basically and 61 reports were shortlisted. In the wake of eliminating copies and rethinking qualifications, three articles were remembered for the last audit. Each of the three investigations remembered for this audit scored scores from 8 to 15 (out of 29) on the danger of pre-

disposition evaluation device. Every one of them scored low on the strength model dependent on the lower number of subjects remembered for these investigations. In detailing and choosing inclination, all examinations scored normal or better than expected. All examinations neglected to accomplish a dumbfounding normal score. Two examinations that showed a positive connection between nutrient D levels and paces of coronavirus disease scored low in the danger evaluation of inclination. An examination that showed no impact of Vitamin D demonstrated a normal score. There is just incidental proof connecting the coronavirus discoveries and nutrient D status. The job of high-portion Nutrient-D against coronavirus should be completely assessed in observational investigations or top-notch randomized examinations before it is suggested.

Since the flare-up of COVID-19, just a piece of the contamination, clinical qualities, and mortality designs have been distinguished. For a pandemic of worldwide extents, we have restricted cure alternatives¹¹. The general death rate detailed in the emergency unit is 25.7%¹². The absence of a conclusive antibody and corrective medication has brought about a combination of medications as analyst's attempt to locate the most fitting cure technique. Accordingly, the assessment of chemoprophylaxis and obstruction techniques is one of the critical methodologies to forestall further disease and mortality. Since the improvement of the antibody will require some serious energy, somewhere in the range of 6 and a year and a half, there is a ton of theory about medications like hydroxychloroquine, nutrient D, nutrient C and zinc¹³⁻¹⁷. News levels and online media have introduced the association of wholesome upgrades inside the remedy and avoidance of Coronavirus Disease 2019 (COVID19). During this pandemic, when information and online media tiers get involved in nutritional enhancements, mixed messages and deception attain all around the vicinity. Nutrient D movement helps in keeping up the cell-actual hindrance, articulation of antimicrobial peptides, the action of macrophages and monocytes, and action of cells engaged with natural and versatile insusceptibility, for example, dendritic cells and T-cells Mechanism^{18,19}. Information from observational investigations interfaces low degrees of nutrient

D with intense respiratory contamination²⁰. Through resistant balance, nutrient D could impact infection replication and help in the insusceptible guideline, and in principle, it could diminish contamination and mortality. The test, in any case, is the deciphering impact of this model on genuine clinical practice. Studies from Italy and different nations feature the utilization of nutrient D, demonstrating that sufficient degrees of nutrient D should be guaranteed through safe openness to the sun, food, or nutrient D supplementation²¹. Our objective was to lead an efficient survey of the writing on the part of nutrient D in coronavirus contamination and mortality, as it isn't realized whether taking nutrient D day by day improves results related to coronavirus².

R3: How can the mortality rate of COVID-19 patients be?

A few ends have been drawn from the demise price of code of federal policies contemplates led with the aid of the USA Public Health Service throughout the 1918-1919 flu pandemic. The code of federal regulations pneumonia rate in the test diary was 28.8% for whites and 39.8% for the "shaded", and Southwestern social orders had a lower code of federal regulations rate than the Northeastern social orders because of dosages; most elevated sun-based UV beams in summer and winter^{22,23}. In a new twofold visually impaired randomized controlled preliminary directed in ventilated ICU patients, the impact of high-portion nutrient D cure on medical clinic stay span, readmission rate, sepsis, and mortality was assessed. Subjects were given ($n = 31$) a false cure, 50,000 worldwide units (IU) of cholecalciferol (nutrient D3), or 100,000 global units of nutrient D3 day by day for 5 successive days (all-out nutrient D3 portion = 250,000 IU or 500,000 units) International, individually). There was a huge decrease long of emergency clinic stay over the long run in the 250,000 IU and 500,000 IU nutrient D3 gathering, contrasted with the false cure gathering (25 + 14 and 18 + 11 days contrasted with 36 + 19 days, individually; $p = 0.03$). Other clinical boundaries didn't show measurably huge contrasts^{24,25}. The significance of remedying nutrient D insufficiency could not be more important in certain networks at high danger for clinic gained diseases, for example, forefront laborers, clinical and medical services laborers, patients and guests. Serum centralizations of 25 (OH) D of in any event 40-50 ng/mL (100-125 nmol/L) are demonstrated based on observational inspections^{3,26,27}.

R4: How can we Investigate the Frequency of Abnormalities Detected by Point-of-Care in the Lung?

The coronavirus pandemic has brought about dreariness and mortality and wellbeing frameworks are underneath stress across the planet. Along those anxieties, techniques are being searched for which could

accelerate and enhance the finding and the board of this element. Recent evidence²⁷ exposed the use of lung ultrasound to diagnose coronavirus by finding an interstitial sequence (type B). Our intention became therefore to pool the percentage of various lung anomalies identified and determined by way of LUS in symptomatic coronavirus sufferers. We performed a systematic assessment (PubMed, MEDLINE, and EMBASE until April 25, 2020) and a pedigree meta-evaluation. Seven experiments investigated lung ultrasound activity in 122 coronavirus sufferers were involved. The pooled ratio (PP) for type B observed by pulmonary ultrasound (US) was 0.97(95% CI 0.94-1.00 I 2 0%, Q 4.6). PP for finding abnormalities in the pleural line was 0.70 (95% CI: 0.13-1.00 I 2 96%, Q 103.9), pleural fusion was 0.54 (95% 0.11-0.95 I 2 93%, Q 61.1) or pulmonary fusion was 0.39 (95% CI: 0.21-0.58 I 2 72%, Q 17.8) and pleural effusion was 0.14 (95% CI 0.00-0.37 I 2 93%, Q 27.3)⁴. Our meta-examination showed that almost all patients with SARS-CoV2 had atypical lung ultrasound filters. The highest documented irregularity is the interstitial addition referred to in Example B. Disclosure of our survey is a workable activity of this technique in the event of an emergency, inference and observation of sufferers of coronavirus. The high indicative accuracy of the comparison tests for lung ultrasound, CT, and COVID-19 is justified by additional consideration of this result and the determination of the symptomatic and prognostic appearance for each of these modalities^{28,29}.

Current attention in factor-of-care ultrasound (POCUS) is because of its portability, steep studying curve, quite less complicated sterilization technique, absence of publicity to ionizing radiation, and its position in tracking. In addition, it has a remarkable dating with CT in various lung sicknesses (B lines, subpleural consolidation and abnormal pluralin). The use of POCUS has a greater fee in aid-restricted settings, as an example, in a few tropical areas in which other diagnostic strategies might not be comfortable to be had and checking outsources can be restrained. In those settings, fundamental talents of ultrasound photo acquisition and interpretation may be taught to fitness care vendors from specific backgrounds via following a brief schooling direction³⁰. In our overview, Type B prevailed, taking place with repeated manipulation (FP) of 97% (94-100%). The Results had been constant and homogeneous throughout all issue research. Pleural line abnormalities had been present in two-thirds of the cases (FP 70%, 95% CI: 1300 %), the frequency of different findings changed into less in our review, and the outcomes have been very heterogeneous. However, this function desires to be supported with the aid of potential studies two research excluded from our assessment are noteworthy^{31,32}. The first take a look at examined the role of lung ultrasound in asymptomatic coronavirus patients and was as a consequence excluded³³. In their retrospective evaluation of 9 asymptomatic patients, LES revealed an abnormality in 22% ($n = 2/9$). One patient had kind B and the opposite patient had lung stiffening at the same

time as the frequency of type B become low. In this study, CT scans likewise confirmed abnormalities in the simplest 33% (n = three/nine), indicating the potential poor overall performance of the exceptional imaging strategies inside the sufferers. Those suffering from coronavirus are asymptomatic. The 2nd study became excluded as it changed into restricted to a group of children. They retrospectively analyzed statistics from eight children with coronavirus: 88% (n = 7/8) had abnormalities on lung ultrasound (type B = 5/8, combo's n = 2/8). In phrases of medical development, all lesions advanced radiologically, either partially or completely, indicating an ability position for ultrasound in medical observe-up. The results of bacterial pneumonia have been pulmonary consolidation using an ultrasound bronchoscopy. However, the effects observed in instances of viral pneumonia were similar to those found for us (kind B, pleural line abnormalities or subpleural fusion)^{34,35}. It may be stated that the LUS outcomes may not permit docs to distinguish among coronavirus and other viral lung infections; however, having any such frequent locating inside the middle of a virus will cause faster diagnostic and Cure selections and better aid allocation. Forest plot represents (a) the mixed ratio of pleural line abnormalities (thickening or irregularity of the pleural, whichever is greater), (b) thickening of the pleura, and (c) pleural effusion detected utilizing lung ultrasound in patients with signs and symptoms of COVID-19 There is a high heterogeneity represented by I2 for the 3 disparities Pulmonary ultrasound and coronavirus. Our assessment aimed to synthesize the proportions stated from the distinctive consequences detected utilizing the lungs of POCUS and is the first meta-analysis aimed toward assessing the function of lung ultrasound, or POCUS, in diagnosing coronavirus. The authors recollect this to be of price to front-line clinicians. We accept as true that the consequences of evaluation will help integrate this beneficial method in triage, prognosis, control and follow-up of patients with coronavirus³⁵.

R5: What are the Most Popular Coronavirus Control Strategies Discussed in the Literature?

The novel coronavirus, due to SARS-CoV-2, has proven one of a kind occurrence and mortality rates (code of federal regulations) amongst sufferers with one-of-a-kind pre-current chronic conditions. Patients with giving up-level renal ailment (ESRT) who are on Renal Replacement Therapy (RRT) may additionally have a higher prevalence and code of federal regulations due to reduced immune feature because of hematuria and renal swelling from SARS-CoV-2; however, there has been no having a look at A methodology on the contamination and mortality of SARS-CoV-2 infection in ESRD patients with RRT. Five electronic databases were searched, and a scientific evaluation and meta-analysis did as of June 30, 2020; to evaluate the prevalence and cause mortality (code of federal regulations) of coronavirus contamination among ERT patients with

RRT. Global coronavirus statistics become retrieved from the global database on June 30, 2020, to estimate the prevalence and code of federal regulations of the overall populace as reference standards. Of the 3, 272 capacity research, 34 eligible kinds of research such as 1944 showed instances of coronavirus have been in 21,873 ERT sufferers with RRT from 12 nations in 4 WHO regions. The combined typical occurrence in ESRD sufferers with RRT became 3.10% [95% confidence interval (CI)], which is above the global reference implying occurrence of 0.14%. The envisioned common code of federal regulations for coronavirus in ESRD patients with an RRT became 18.06% (95% CI: 14.09-46-22.32), better than the global common of 4.98%. This meta-evaluation counseled a higher incidence of coronavirus and code of federal regulations in sufferers with ESRD and RRT. Patients with ESRD and RRT must have the coronavirus Preclusion and Cure Protocol to relieve extra cases and deaths¹⁰.

Methods

The proposed analysis includes the statistical compilation of data related to retroactive trials in a chain of computed tomography experiments from around the world. The findings confirmed an advanced standard for the identity of atypical statistical records from coronavirus sufferers with computed tomography traits. Sampling training is used for coronavirus and coronavirus research. It has been turned into a fantastic detection device around the world. A multicentre retrospective explores how well the variation of coronavirus wall detection walls in China has established that this version can differentiate coronavirus from network-obtained pneumonia (NOP). To affirm the present situation of the general arena, we administer the statistical outcomes of the coronavirus-affected person log to check the modern state of affairs of the general international.

Sampling Techniques

Sampling strategies are classically obtained from accessible data sources applicable to coronavirus, ranging from numerical data on infectious events, radiological images, natural languages to biological data. These results are useful for scientific purposes to harness the potential and strength of medical imaging technology in the tough fight in contradiction of fatal coronavirus disease.

Sampling proof is based on human saliva and lung experiment images. The pattern length is extra than 5000 target population for the harm due to the cases of coronavirus and the human sample collection. The records become entered and evaluated the usage of the new version 2.6 of the SPSS. Most of them had been satisfied with the brand new COVID-19 final results method. The key regions of attention had been the importance of statistical facts, time manipulation and recognition of the primary area of the affected person difficulty.

Hypothesis

R1: What is the overall communal death rate for pandemic coronavirus patients?

R2: What are the effects of deficiency of Vitamin D for pandemic coronavirus?

R3: How can be Mortality rate of COVID-19 Patients?

R4: How can we investigate the Abnormalities Frequency of Detected by the cure of Lungs?

R5: What is the most popular coronavirus control strategies discussed in the literature?

Search Approaches

Before the systematic study of mapping was published, the “empirical research on Coronavirus” series was presented in Science Direct³⁶. In this informal search method, several observational analyses have been identified, so that a scientific mapping appearance has been accepted, and the preliminary set, which has been updated to a prescribed search strategy, has become the statistical results of coronavirus. When this tentative string changed into blanketed inside the Science Direct search engine, the returns obtained no longer consisted of all the numbers at the search engine. In addition, senior software program engineering pupils from lecturers who recognize SLRs have been selected as professionals and had been invited to review the quest string and offer enter. Based on these expert analyses, the original string turned up to date and the number one seek string was separated into components. Expert opinion is a manner to without problems compare and verify records³⁷. Below are the two factors of our described series.

Academic Databases

In this research, we use the following database for the search engine of this mapping of coronavirus cases to be used the digital libraries are given below:

- Embase
- Ovid Medline
- Ovid Cumulative
- Science Direct
- Wiley

The above databases have been determined as common venues for publishing AI-coronavirus issues. Other scholars have used these records extensively in their SLR research^{35,38,39}.

Periodical Collection

This paper details the inclusion and exclusion criteria required for the probability of publishing and also reveals the process used to choose the required courses according to the questions of the studies. The following conditions for inclusion have been laid down: The appraisal period has changed to almost ten years and

consists of studies from 2019 to 2020. This start date is selected because the highest coronavirus (coronal diseases) was reported in 2019 and after. However, the search was carried out at the beginning of 2019, so that the simplest courses starting at the end of 2018 were included in the structured mapping of a look.

- Observational study with a coronavirus emphasis.
- Experiments that consider the coronavirus. The following exclusion criteria have been used:
- Reports that do not contain clear facts about how to keep a coronavirus issue file and medical data.
- Studies of which results are not empirically tested.
- Research is accessible only as abstracts or Power-Point presentations.
- Articles containing the most appropriate instructions, guidelines or sketches of the Coronavirus precaution.
- Introductory articles on seminars, special issues and novels.
- Book chapters, please.
- Articles that are no longer printed in English.

The process for selecting courses has become automated and duplicate: first, the initial selection from the search result has been rendered in compliance with the selection criterion by using the title and abstract of the guides; second, the papers chosen during the preliminary phase have been thoroughly read if you wish to pick the guides for the very last selection, based entirely on the definition of presence criteria.

In this fast-growing field of digital engineering, it's not as straightforward as it was before, because of the strong demands of the consumer and the outstanding nature of the goods. In reality, at this stage of globalization, it will be impossible to properly plan the project, complete it on demand and make it possible to succeed on time with minimal resources. On this statistic, the review of the research is made on certain SPSS software for the results of the overall knowledge for COVI-19 statistically to be considered for the high quality and effectiveness of the challenge. In information engineering, an adaptive approach has been put in place to test meaningful outcomes to grasp the overall scenario of the complete case file in the Coronavirus archive from December 2019 to March 2021 to obtain successful results.

RESULTS

The subsequent causes are used to assess this study which is given below:

The Tables 1, 2 and 3 mentioned the details of publishing databases and the selection of manuscripts in this research. Libraries, including Embase, Ovid Cumulative and Science Direct, extracted courses related to the field of cutting-edge, have all been published in journals. Out of these three libraries, the frequency of the Research Direct book has shifted to the very highest of 25% of the papers in the pool Embase and Ovid Cumulative were second and third in the selected, respectively. Table 4 displays the overall not uncommon positions for studies number one with a frequency of two or more. The findings show that most of the papers are written in journals.

Table 1. Study selection.

Source	Retrieved	Initial selection	Final selection
<i>Embase</i>	102	10	10
<i>Ovid Medline</i>	80	20	3
<i>Ovid Cumulative</i>	50	50	8
<i>Science Direct</i>	50	20	3
<i>Wiley</i>	40	15	5
Total	322	115	29

Table 2. Distribution of studies.

	Journal Manuscript	Conference Manuscript	Workshop Manuscript	Total
<i>Embase</i>	7	2	1	10
<i>Ovid Medline</i>	2	1	0	3
<i>Ovid Cumulative</i>	7	0	1	8
<i>Science Direct</i>	3	0	0	3
<i>Wiley</i>	4	1	0	5
Total	23	4	2	29

Table 3. Selected studies.

Library	Type	Frequency
<i>Embase</i>	Journal	5
<i>Ovid Medline</i>	Journal	2
<i>Ovid Cumulative</i>	Workshop	1
<i>Science Direct</i>	Conference	1
<i>Wiley</i>	Journal	1

spectrum and length of the GGO and a gradual development of the GGO into a stable multifocal opacity, a septal defect, and an insane paving sequence, with the best CT results obtained on the tenth day after the onset of signs and symptoms. Acute respiratory distress syndrome is the most common reason that coronavirus patients are admitted to the intensive care unit and is the leading cause of death of these patients⁴⁰⁻⁴⁸. Clinical developmental imaging trends normally occur after the second week of illness and contain the sluggish resolution of stable opacities and loss of lesions and damaged lobes. This comprehensive analysis of recent coronavirus literature includes statistics on the basic and controlled characteristics of CT disease⁴⁹. On 7 January 2020, the unconventional coronavirus (2019-nCoV; later renamed SARSCoV-2) turned into showed as the motive of the recorded instances, and the epidemic changed into later called the “Coronavirus”⁵⁰⁻⁵⁵.

DISCUSSION

Refereeing to R1:

This manuscript addresses a variety of reporting research databases such as PubMed, Embase (Elsevier), Google Scholar, and WHO databases. To Follow-up CT images at the core stage of the disease demonstrate an increase in the

Table 4. Statistical result of the total number of patients records.

Statistics	Date	Day	Month	Year	Record Cases	Deaths	
Total Number	Valid	25935	25935	25935	25935	25935	
	Missing	0	0	0	0	0	
Mean		15.92	4.15	2020.00	369.45	18.86	
Std. Error of Mean		0.054	0.010	0.000	13.009	0.758	
Median			16.00	4.00	2020.00	4.00	0.00
Mode			26	5	2020	0	0
Std. Deviation			8.653	1.548	0.051	2094.975	122.095
Variance			74.875	2.398	0.003	4388918.684	14907.092
Range			30	11	1	57232	6846
Minimum			1	1	2019	-2461	-1918
Maximum			31	12	2020	54771	4928
Sum			412962	107660	52388633	9581803	489182
Percentiles	25		9.00	3.00	2020.00	0.00	0.00
	50		16.00	4.00	2020.00	4.00	0.00
	75		23.00	5.00	2020.00	65.00	1.00

Chest radiographs have no diagnostic value in the early stages, even though the results of CT experiments can be a blessing even before signs and symptoms appear^{56,57}. In the middle to advanced levels of the sickness, chest X-rays can display growth inside the characteristics of acute respiratory misery syndrome (ARDS). In addition, CT outcomes were proven to be diagnostic in many instances with a false lousy preliminary RTPCR assay^{58,59}. In all cases, an examination of 41 patients with a laboratory-shown diagnosis of coronavirus shows remarkable symptoms of chest CT⁶⁰. Next, a couple of case collections were released to symbolize the coronavirus pneumonia CT scans. With the growing global problem around the outbreak of coronavirus, comprehensive knowledge on diagnostic imaging trademarks, unusual features, and the advancement of chest imaging findings are necessary for the power management and remedy of the affected Individual. However, the material is dispersed in a variety of courses and there has not yet been a full review of the literature⁶⁰.

Refereeing to R2:

Our database seeks returned fifty-nine data and a manual seek of associated articles, and previous systematic critiques delivered two more statistics. After eliminating the duplicates, fifty-nine statistics have been diagnosed for the first step inside the choice manner. From the initial display screen for titles and abstracts, fifty-one facts have been excluded. For the closure of the 5 publications protected by the full-text assessment, three articles were sooner or later omitted. No randomized controlled trials were found among the 3 papers that met the inclusion criterion. One observes was retrospective, the alternative was retrospective, and the alternative is an evaluation⁶¹⁻⁶³. All 3 researchers covered in this evaluation scored scores of 8 to 15 (out of 29) at the risk of the prejudice assessment unit. Many of them ranked poor on the energy criteria based on the lower quantity of topics covered by this study. All studies ranked average or above common in reporting and decision-making on the bias.

Refereeing to R3:

There is most effective circumstantial evidence linking coronavirus outcomes to diet D popularity, and free of nutrition D trying out and subsequent supplementation is a large economic burden for most people in devel-

oping international locations. The function of a Journal Pre-proof 10 high-dose nutrition D towards Coronavirus has to be evaluated in observational research or exceptional randomized studies earlier than recommending it.

Table 4 shows the statistical result of the total number of patients records such as death cases, case report legally, year with month and date. We collect this data online overall the world as Google has compiled online the information base created, installed and proven with the aid of the World Health Organization and the Centers for Disease Control and Disease Preclusion (ECDP) for almost all health center cases filed due to coronavirus. The records changed into accumulated in some unspecified time in the future of the length 2019-2020 and the ensuing responses have been feed into the SPSS “Statistics Package for Computational Research” for further examination. In the SPSS, the researcher makes use of the *t*-test a look at the statistics obtained for assessment. In this evaluation, statistics are evaluated by way of SPSS. SPSS is a statistical technique broadly utilized by many researchers. The *t*-test shows the significant ratio which is used to evaluate the affiliation between the CT experiment opinions of the coronavirus patient sequences as proven by Figures 3 and 4.

Figures 3 and 4 show the statistical results of death cases of COVID-19 patients overall the world in a monthly, yearly and weekly report with countries and territories. The above figure mentions the cases of most effective countries due to COVID-19 from the beginning to 2020-year scenario.

Refereeing to R4:

Our overview is restrained by way of a small quantity of problem research, a small pattern of patients, a borderline bias vulnerability and incapacity to get rid of bias from e-books. Furthermore, there may be a scarcity of steady definitions and a loss of accuracy within the documentation of certain lung abnormalities. Insufficient documentation of the significance of the LUS final results (affected lung areas or representative LUS rating) can restrict its function in temporal enforcement and predictive fee. At the quilt of the day, we have been not able to quantify the sensitivity and accuracy of the missed information for their account. Effective diagnostic accuracy because of LUS analysis, CT scans, and numerous coronavirus (PCR, IgM, and IgG) tests of serial measurements to de-




Figure 3. Statistical results of death cases of COVID-19 patients overall the world.




Figure 4. Countries and territory representation of COVID-19 cases.

cide the sensitivity, specificity and accuracy of the coronavirus diagnostic approach is crucial. In addition, LUS analysis in sickness-precise depth spectra is proposed to degree effects that can be related to disease severity. The outcomes of the maximum current and ongoing research, together with POCUSCO, ECHOVID-19, POCUSars-CoV-2, VIRUS and COVILUS, would deal with a number of the shortcomings set out above^{64,65}.

Refereeing to R5:

Famous code of government guidelines of COVID-19 in End-Stage Renal Disease patients expanded to higher than code of administrative guidelines around the world (18.06% vs. 3574.98%)³. Conversely, the surmised normal code of government guidelines of COVID-19 in ESRD patients was higher than in everybody (18.06% vs. 0.6% vs. 8.5%)⁶⁶, hypertension (18.06% vs. 6.4%) and immune system issues (18.06% vs. 6.60%), however falls because of diabetes (18.06% vs. 19.25%)⁷⁶ and malignancy patients (18.06% vs. 20%). Disabled resistant capacity can assume a critical part in the extension of the code of government guidelines in ERT patients with RRT⁷⁷. Notwithstanding no proof of characteristic inclination, there has been significant fluctuation between WHO-level case-casualty preliminaries from 9.51% in the western Pacific to 17.01% in the Americas and 18.77% in the Americas and European territories. The code of government guidelines adaptation might have corresponded with the abnormally low rate of COVID-19 broadly in the western Pacific zone of China comparative with other included nations⁷⁵. A few lacks ought to be found in this orderly examination and meta-assessment. Second, the included example showed enormous varieties in both pervasiveness and RFC results. The subgroup research did not altogether investigate all wellsprings of heterogeneity, considering the incomplete accessibility of information in the included examinations. Second, there was no sound test for ESRD patients with worked on PD⁶⁷⁻⁷⁰. The pervasiveness of COVID-19 and the code of government guidelines in PD patients would likewise not be unavailable. Third, the specifics of the TSR process, including HD

type, KT type, and TSR length, were also not discussed. Finally, we cover how-to posts in English that can be read by the facts pages in various languages.

Finally, our systematic analysis and meta-evaluation presented the first proof of co-occurrence and coronavirus code of federal regulations in ESRD patients with RRTs above the global average. Increased prevalence and mortality may be due to decreased immune function in patients with End-Stage Renal Disease⁷¹⁻⁷⁵. ESRD patients with RRT should have their regimen for the detection and care of coronavirus for the alleviation of excess cases and deaths⁷⁶.

CONCLUSIONS

The goal of this research is to offer an outline of the Coronavirus disorder that has seriously affected the lives of people around the sector. Many human beings lost loved ones; the death instances global exceed three hundred million and keep to upward push. There is insufficient evidence to be had to illustrate the accuracy of the CT or deep gaining knowledge of the model that makes use of detection networks to detect/locate sufferers with COVID-19 among the ones without COVID-19 or community-acquired pneumonia. For decades, many vaccines, such as the measles or polio vaccines, were protective but disappeared over time, such as whooping cough and influenza. Long-term immunity is also caused by some forms of viral infection and others by a transient response. Here are some of the announced results of the systematic reviews of COVID-19 listed below:

R1: This systematic evaluation becomes collected to establish the functions of CT scans of this rapidly spreading and life-threatening viral pneumonia, to summarize key imaging observations at various ranges of the sickness, and to determine the progression of CT findings steady with ailment development or clinical recuperation.

R2: There is only circumstantial proof concerning the effects of coronavirus on the repute of nutrition D. coronavirus seemed and dispersed because the

Northern Hemisphere witnessed winter (past due 2019) and 25-hydroxyvitamin D levels peaked at the lowest stage. There is also an excessive range of instances and deaths in those areas. While it could be visible that countries consisting of Norway, Sweden and Finland, which are lighter than southern Europe, have a decrease occurrence and mortality rate, their population have a far better average of 25 (OH) D and are therefore comparatively enough nutrition D due to massive-scale fortification ingredients. On the alternative hand, Italy and Spain, international locations of decrease range, are both exceptions; however, the prevalence of diet D deficiency in these populations is remarkably commonplace. Similarly, it seems that darkish-skinned races inclusive of blacks and ethnic minorities who are more susceptible to Vitamin-D deficiency due to reduced absorption of Ultraviolet-B (UVB) rays tend to be greater affected than those with fair pores and skin. A currently posted look explores the function of vitamin D in the preclusion of influenza and the way supplementation may be a useful measure to lessen the risk of developing COVID-19.

R3: This model suggests the connection between the established and independent variable that the output is interpreted. As an impact of two-length of the cases, there might be a disparity between the ways; however, the route of the distinction cannot be expected.

R4: Proof of interstitial lung activity, as visible in the B-pattern, is the maximum common and reliable finding at the US lung in coronavirus patients. While non-specific the prevalence of this finding amid coronavirus pandemic, further to other feature signs, would boom the threat of contamination. Thus, POCUS is probable to play a critical position within the capability triage, analysis, cure and follow-up of coronavirus sufferers.

R5: In addition, cure plans together with antiviral marketers, vaccines and oxygen therapy which can be utilized in extra than 3 quarters of patients are now being analyzed. The findings could have a “proof-primarily based wish” about how to deal with this rapid and daunting epidemic.

In the end, our Meta-data analysis and systematic evaluation process obtainable the proof of occurrence and code of federal regulations of coronavirus that become the developed than the global common in ESRD patients with RRT.

Increased incidence and mortality may be related to reducing immune characteristics in patients with ESRD. ESRD sufferers with RRT need to offer their own coronavirus avoidance and recovery regimen to lessen excess cases and deaths.

COMPETING INTERESTS:

Dr. Soobia Saeed and Prof. Dr. A.S. Hashim are working on Neurospinal and Cancer Care Institute and Hospital, Karachi, Pakistan. They are recently working on COVID-19 patients and deal with chronic cases for the treatment of Breathing issues and frequency develop the abnormalities cases created by Coronavirus. Other authors declare no conflict of interest.

CONFLICT OF INTEREST:

The authors declare that they have no conflict of interest.

REFERENCES

- Hatmi ZN. A Systematic Review of Systematic Reviews on the coronavirus Pandemic. *SN Compr Clin Med* 2021; 26: 1-8.
- Yadav SK, Gaurav K, Johri G, Jaiswal SK, Jha CK, Yadav N. A systematic review of the role of hypovitaminosis-D in coronavirus disease-19 (coronavirus) infection and mortality: Is there a role of recommending high dose Vitamin-D supplementation. *Human Nutrition & Metabolism* 2021; 28: 200120.
- Mohamed MF, Al-Shokri S, Yousaf Z, Danjuma M, Parambil J, Mohamed S, Mubasher M, Dauleh MM, Hasanain B, AlKahlout MA, Abubeker IY. Frequency of abnormalities detected by point-of-care lung ultrasound in symptomatic Coronavirus patients: systematic review and meta-analysis. *Am J Trop Med Hyg* 2020; 103: 815-821.
- Milovanovic L, Hessey E, Sebastianski M, Keto-Lambert D, Vandermeer B, Bagshaw SM, Rewa O. Epidemiology Clinical characteristics and Cure of critically ill patients with coronavirus: a protocol for a living systematic review. *BMJ* 2021; 11: 1-5.
- Karlsen AP, Wiberg S, Laigaard J, Pedersen C, Rokamp KZ, Mathiesen O. A systematic review of trial registry entries for randomized clinical trials investigating coronavirus medical preclusion and Cure. *PLoS One* 2020; 15: 1-13.
- Meyerowitz EA, Sen P, Schoenfeld SR, Neilan TG, Frigault MJ, Stone JH, Kim AY, Mansour MK. Immunomodulation as Cure for Severe coronavirus: a systematic review of current modalities and future directions. *Clin Infect Dis* 2020; 1-14.
- Nopsopon T, Kittrakulrat J, Takkavatakarn K, Eiamsittrakoon T, Kanjanabuch T, Pongpirul K. Coronavirus in end-stage renal disease patients with renal replacement therapies: A systematic review and meta-analysis. *MedRxiv* 2021; 77-105.
- Salehi S, Abedi A, Balakrishnan S, Gholamrezaezhad A. Coronavirus disease 2019 (coronavirus): a systematic review of imaging findings in 919 patients. *AJR Am J Roentgenol* 2020; 215: 87-93.
- Ghayda RA, Lee J, Lee JY, Kim DK, Lee KH, Hong SH, Han YJ, Kim JS, Yang JW, Kronbichler A, Smith L. Correlations of clinical and laboratory characteristics of coronavirus: A systematic review and meta-analysis. *Int J Environ Res Public Health* 2020; 14: 5026.
- Nopsopon T, Kittrakulrat J, Takkavatakarn K, Eiamsittrakoon T, Kanjanabuch T, Pongpirul K. Coronavirus in end-stage renal disease patients with renal replacement therapies: A systematic review and meta-analysis. *MedRxiv* 2021; 1-12.
- Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, Ruan L, Song B, Cai Y, Wei M, Li XW, Xia J, Chen N, Xiang J, Yu T, Bai T, Xie X, Zhang L, Li C, Yuan Y, Chen H, Li H, Huang H, Tu S, Gong F, Liu Y, Wei Y, Dong C, Zhou F, Gu X, Xu J, Liu Z, Zhang Y, Li H, Shang L, Wang K, Li K, Zhou X, Dong X, Qu Z, Lu S, Hu X, Ruan S, Luo S, Wu J, Peng L, Cheng F, Pan L, Zou J, Jia C, Wang J, Liu X, Wang S, Wu X, Ge Q, He J, Zhan H, Qiu F, Guo L, Huang C, Jaki T, Frederick G, Hayden, Peter W, Horby, Zhang D, Wang C. A trial of lopinavir-ritonavir in adults hospitalized with severe Covid-19. *N Engl J Med* 2020 ;382: 1787-1799.
- Quah P, Li A, Phua J. Mortality rates of patients with Coronavirus in the intensive care unit: a systematic review of the emerging literature. *Crit Care* 2020; 24: 285.
- Gao J, Tian Z, Yang X. Breakthrough: chloroquine phosphate has shown apparent efficacy in the Cure of Coronavirus associated pneumonia in clinical studies. *BioSci Trends* 2020; 14: 72-73.

14. Gautret P, Lagier JC, Parola P, Hoang V, Meddeb L, Mailhe M, Doudier B, Courjon J, Giordanengo J, Vieira VE, Dupont HT, Honoré S, Colson P, Chabrière E, Scola BL, Rolain JM, Brouqui P, Raoult D. Hydroxychloroquine and azithromycin as a Cure of Coronavirus: results of an open-label non-randomized clinical trial. *International Journal of Antimicrobial Agents* 2020; 20:105949.
15. Shah S, Das S, Jain A, Misra DP, Negi VS. A systematic review of the prophylactic role of chloroquine and hydroxychloroquine in coronavirus disease-19 (Coronavirus). *Int J Rheum Dis* 2020; 23: 613-619.
16. Kara M, Ekiz T, Ricci V, Kara Ö, Chang KV, Özçakar L. Scientific Strabismus or Two Related Pandemics: Coronavirus & Vitamin D Deficiency. *Br J Nutr* 2020; 1-20.
17. Adams KK, Baker WL, Sobieraj DM. Myth Busters: Dietary Supplements and coronavirus. *Annals of Pharmacotherapy* 2020; 54: 820-826.
18. Prietl B, Treiber G, Pieber TR, Amrein K. Vitamin D and immune function. *Nutrients* 2013; 5: 2502-2521.
19. Aranow C. Vitamin D and the immune system. *J Investig Med* 2011; 59: 881-886.
20. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, Bhattoa HP. Evidence that vitamin D supplementation could reduce risk of influenza and coronavirus infections and deaths. *Nutrients* 2020; 988: 2-19.
21. Lau FH, Majumder, R, Torabi, R, Saeg, F, Hoffman, R, Cirillo JD, Greiffenstein P. Vitamin D insufficiency is prevalent in severe COVID-19. *MedRxiv* 2020; 1-14.
22. Ilie PC, Stefanescu S, Smith L. The role of vitamin D in the preclusion of coronavirus disease 2019 infection and mortality. *Aging Clin Exp Res* 2020; 1-4.
23. Grant WB, Giovannucci E. The possible roles of solar Ultraviolet-B radiation and vitamin D in reducing case-fatality rates from the 1918–1919 influenza pandemic in the United States. *Dermatoendocrinol* 2009; 1: 215-219.
24. Feigenbaum, Muller JJ. Regional and Racial Inequality in Infectious Disease Mortality in U.S. Cities, 1900–1948. *Demography* 2019; 56: 1371-1388.
25. Han, JE, Jones JL, Tangpricha V, Brown, Hao L, Hebbar G Lee, MJ Liu, S Ziegler. High Dose Vitamin D Administration in Ventilated Intensive Care Unit Patients: A Pilot Double-Blind Randomized Controlled Trial. *J Clin Transl Endocrinol* 2016; 4: 59-65.
26. Smith EM, Jones JL, Han JE, Alvarez JA, Sloan JH, Konrad RJ, Tangpricha V. High-dose vitamin D3 administration is associated with increases in hemoglobin concentrations in mechanically ventilated critically ill adults: a pilot double-blind, randomized, placebo-controlled trial. *J Parenter Enteral Nutr* 2018; 42: 87-94.
27. Youssef, Ranasinghe T, Grant WB Peiris AN. Vitamin D's potential to reduce the risk of hospital-acquired infections. *Dermatoendocrinol* 2012; 4: 167-175.
28. Howie SEM, Norval M, Maingay J. Exposure to low dose UVB light suppresses delayed-type hypersensitivity to herpes simplex virus in mice. *J Invest Dermatol* 1986; 86: 125-128.
29. Halliday GM. Inflammation, gene mutation and photoimmunosuppression in response to UVR-induced oxidative damage contributes to photocarcinogenesis. *Mutat Res* 2005; 571: 107-120.
30. Wang L, Jameson SC, Hogquist KA. Epidermal Langerhans cells are not required for UV-induced immunosuppression. *J Immunol* 2009; 183: 5548-5553.
31. Grewe M, Gyufko K, Krutmann J. Interleukin-10 production by cultured human keratinocytes: regulation by ultraviolet B and ultraviolet A1 radiation. *J Invest Dermatol* 1995; 104: 3-6.
32. Terpos E, Ntanasis-Stathopoulos I, Elalamy I, Kastritis E, Sergeantanis TN, Politou M, Dimopoulos MA. Hematological findings and complications of COVID-19. *Am J Hematol* 2020; 95: 834-847.
33. Kinlen LJ, Sheil AG, Peto J, Doll R. Collaborative United Kingdom-Australasian study of cancer in patients treated with immunosuppressive drugs. *Br Med J* 1979; 2: 1461-1466.
34. Hamzavi IH, Lyons AB, Kohli I. Ultraviolet germicidal irradiation: Possible method for respiratory disinfection to facilitate reuse during the coronavirus pandemic. *J Am Acad Dermatol* 2020; 82: 1511-1512.
35. Totonchy MB, Chiu MW. UV-based therapy. *Dermatol Clin* 2014; 32: 399-413.
36. Michelle A Bylicky, Gregory P Mueller, Regina M Day. Radiation resistance of normal human astrocytes: the role of non-homologous end-joining DNA repair activity. *J Radiat Res* 2019; 60: 37-50.
37. Fried JA, Ramasubbu K, Bhatt R, Topkara VK, Clerkin KJ, Horn E, Rabbani L, Brodie D, Jain SS, Kirtane AJ, Masoumi A, Takeda K, Kumaraiah D, Burkhoff D, Leon M, Schwartz A, Uriel N, Sayer G. The Variety of Cardiovascular Presentations of COVID-19. *Circulation* 2020; 141: 1930-1936.
38. Akhmerov A, Marbán E. COVID-19 and the Heart. *Circ Res* 2020; 126: 1443-1455.
39. Pellicori P. At the heart of COVID-19. *Eur Heart J* 2020; 41: 1830-1832.
40. Wang C, Horby PW, Hayden FG. A novel coronavirus outbreak of global health concern. *Lancet* 2020; 395: 470-473.
41. Phan LT, Nguyen TV, Luong QC. Importation and human-to-human transmission of a novel coronavirus in Vietnam. *N Engl J Med* 2020; 382: 872-874.
42. Riou J, Althaus CL. The pattern of early human-to-human transmission of Wuhan 2019 novel coronavirus (2019-nCoV). *Euro Surveill* 2020; 25: 20-29.
43. Wu Z, McGoogan JM. Characteristics of and Important Lessons from the Coronavirus Disease 2019 (Coronavirus) Outbreak in China: Summary of a Report of 72 314 Cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020; 323: 1239-1242.
44. Huang C, Wang Y, Li X. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; 395: 497-506.
45. Grasselli G, Zangrillo A, Zanella A. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region. *JAMA* 2020; 323: 1574-1581.
46. Grasselli G, Pesenti A, Cecconi M. Critical care utilization for the coronavirus outbreak in Lombardy, Italy: early experience and forecast during an emergency response. *JAMA* 2020; 323: 1545-1546.
47. Moghadas SM, Shoukat A, Fitzpatrick MC. Projecting Hospital utilization during the Coronavirus outbreaks in the United States. *Proc Natl Acad Sci U S A* 2020; 117: 9122-9126.
48. Du RH, Liu LM, Yin W. Hospitalization and critical care of 109 decedents with Coronavirus pneumonia in Wuhan, China. *Ann Thorac Surg* 2020; 17: 839-846.
49. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J. Grade guidelines: Introduction-Grade evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011; 64: 383-394.
50. Waheed A, Sly WS. Carbonic anhydrase XII functions in health and disease. *Gene* 2017; 623: 33-40.
51. Lei Wu, Xuesi Zhou, Zhigang Xiao, Xiujie Gao, Ziquan Liu, Zhiqing Zhang, Kun Wang, Yulin Zhu, Hanlin Ren, Tianhui Wang. Functional expression, characterization, and application of human S100B. *Oncol Rep* 2017; 38: 2309-2316.
52. Karolina Orywal, Maciej Szmikowski, Alcohol dehydrogenase and aldehyde dehydrogenase in malignant neoplasms. *Int J Clin Exp Med* 2017; 17: 131-139.
53. Foroutan F, Guyatt G, Zuk V, Vandvik PO, Alba AC, Mustafa R. Grade guidelines 28: use of Grade for the assessment of evidence about prognostic factors: rating certainty in identification of groups of patients with different absolute risks. *J Clin Epidemiol* 2020; 121: 62-70.

54. Guyatt GH, Ebrahim S, Alonso-Coello P, Johnston BC, Mathioudakis AG, Briel M. Grade guidelines 17: assessing the risk of bias associated with missing participant outcome data in a body of evidence. *J Clin Epidemiol* 2017; 87: 14-22.
55. Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coello P, Rind D. GRADE guidelines 6. Rating the quality of evidence— imprecision. *J Clin Epidemiol* 2011; 64: 1283-1293.
56. Wilder-Smith A, Freedman. Isolation, quarantine, and social distancing and community containment: a pivotal role for old-style public health processes in the novel coronavirus (2019-nCoV) outbreak. *Int J Travel Med Glob Health* 2020; 27: 20.
57. Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M. Grade guidelines: 7. Rating the quality of evidence— inconsistency. *J Clin Epidemiol* 2011; 64: 1294-1302.
58. Ilie PC, Stefanescu S, Smith L. The role of vitamin D in the preclusion of coronavirus disease 2019 infection and mortality. *Aging Clin Exp Res* 2020; 1-4.
59. Hastie CE, Mackay DF, Ho F, Celis-Morales CA, Kattikireddi SV, Niedzwiedz CL, Jani BD, Welsh P, Mair FS, Gray SR, O'Donnell CA, Gill JM, Sattar N, Pell JP. Vitamin D concentrations and coronavirus infection in UK Biobank. *Diabetes Metab Syndr* 2020; 14: 561-565.
60. D'Avolio A, Avataneo V, Manca A, Cusato J, Nicolò, Renzo AD Lucchini, Keller F, and Cantù M. 25-Hydroxyvitamin D Concentrations Are Lower in Patients with Positive PCR for SARS-CoV-2. *Nutrients* 2020; 9: 1359-1362.
61. Loeb M, Dang AD, Thiem VD, Thanabalan V, Wang B, Nguyen NB, Tran HTM., Luong TM., Singh P, Smieja M. Effect of Vitamin D supplementation to reduce respiratory infections in children and adolescents in Vietnam: A randomized controlled trial. *Influenza Other Respir Viruses* 2019; 13: 176-183.
62. Urashima M, Segawa T, Okazaki M, Kurihara M., Wada Y, Ida H. Randomized trial of vitamin D supplementation to prevent seasonal influenza A in schoolchildren. *Am J Clin Nutr* 2010; 91: 1255-1260.
63. Zhou J, Du J, Huang L, Wang Y, Shi Y, Lin H. Preventive Effects of Vitamin D on Seasonal Influenza A in Infants: A Multicenter, Randomized, Open, Controlled Clinical Trial. *Pediatric Infect Dis* 2018; 37: 749-754.
64. Kermali M, Khalsa RK, Pillai K, Ismail Z, Harky A. The role of biomarkers in the diagnosis of coronavirus - a systematic review. *Life Sci* 2020; 254: 117788.
65. Jain V, Yuan JM. Predictive symptoms and comorbidities for severe coronavirus and intensive care unit admission: a systematic review and meta-analysis. *Int J Public Health* 2020; 65: 533-546.
67. Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M. Grade guidelines: 8. Rating the quality of evidence— indirectness. *J Clin Epidemiol* 2011; 64: 1303-1310.
68. Pacifico A, Ardigò M, Francione P, Damiani G, Morrone A. Phototherapeutic approach to dermatology patients during the 2019 coronavirus pandemic: real-life data from the Italian red zone. *Br J Dermatol* 2020; 183: 375-376.
69. Wong LC, Jamil A, Md nor N. The effect of pre-photo test sun exposure on minimal erythema dose and minimal melanogenic dose among skin phototypes III, IV and V. *Photodermatol Photoimmunol Photomed* 2018; 34: 400-404.
70. Blakely K, Drucker A, and Rosen C. Drug-Induced Photosensitivity—An Update: Culprit Drugs, Preclusion and Management. *Drug Safety* 2020; 827-847
71. Rai R, Thomas M. Photopatch and UV-irradiated patch testing in photosensitive dermatitis. *Indian Dermatol Online J* 2016; 7: 12-16.
72. Fujimoto N, Danno K, Wakabayashi M. Photosensitivity with eosinophilia due to ambroxol and UVB. *Contact Dermatitis* 2009; 60: 110-113.
73. Chowdhury MS, Rathod J, Gernsheimer J. A rapid systematic review of clinical trials utilizing chloroquine and Hydroxychloroquine as a Cure for Coronavirus. *Acad Emerg Med* 2020; 27: 493-504
74. Das S, Bhowmick S, Tiwari S, Sen S. An updated systematic review of the therapeutic role of hydroxychloroquine in coronavirus disease-19 (coronavirus). *Clin Drug Investig* 2020; 40: 591-601.
75. Jankelson L, Karam G, Becker ML, Chinitz LA, Tsai MC. QT prolongation, torsades de pointes, and sudden death with short courses of chloroquine or hydroxychloroquine as used in COVID19: a systematic review. *Heart Rhythm* 2020; 17: 1472-1479.
76. Huang I, Pranata R. Lymphopenia in severe coronavirus disease2019 (coronavirus): systematic review and meta-analysis. *J Intensive Care* 2020; 8: 36.